


(9)



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Bc

Search PubMed

for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

Show:

20

Sort

Send to

Text

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

☐ 1: Mol Pharmacol. 2000 Jun;57(6):1165-72.

Related Articles, Links

FREE full-text article of  
www.molpharm.org

## Two different signaling mechanisms involved in the excitation of rat sympathetic neurons by uridine nucleotides.

**Bofill-Cardona E, Vartian N, Nanoff C, Freissmuth M, Boehm S.**

Department of Pharmacology, University of Vienna, Austria.

UTP stimulates transmitter release and inhibits M-type K(+) channels in rat superior cervical ganglion neurons via G protein-coupled P2Y receptors. To investigate the underlying signaling mechanisms, we treated the neurons with either pertussis or cholera toxin; neither treatment altered the inhibition of M-type K(+) channels by 10 microM UTP. However, pertussis toxin reduced UTP-evoked [(3)H]noradrenaline release by 66%. UTP, UDP, ATP, and ADP caused accumulation of inositol trisphosphate in a pertussis toxin-insensitive manner. Pharmacological inhibition of inositol trisphosphate-induced Ca(2+) release (by inhibition of phospholipase C, of inositol trisphosphate receptors, and of the endoplasmic Ca(2+)-ATPase) prevented the UTP-dependent inhibition of M currents but failed to alter UTP-evoked [(3)H]noradrenaline release. Chelation of intracellular Ca(2+) by 1,2-bis(2-aminophenoxy)ethane-N, N,N',N'-tetraacetic acid also reduced the inhibition of M currents by UTP. In addition, all these manipulations attenuated the inhibition of M currents by bradykinin, but hardly affected the inhibitory action of oxotremorine M. These results demonstrate that UTP inhibits M-type K(+) channels via an inositol trisphosphate-dependent signaling cascade that is also used by bradykinin but not by muscarinic acetylcholine receptors. In contrast, the secretagogue action of UTP is largely independent of this signaling cascade but involves pertussis toxin-sensitive G proteins. Thus, UTP-sensitive P2Y receptors excite sympathetic neurons via at least two different signal transduction mechanisms.

PMID: 10825387 [PubMed - indexed for MEDLINE]

Display

Abstract

Show:

20

Sort

Send to

Text